arXiv:1205.2586v2 [gr-qc] 24 Jun 2012

QUANTUM GRAVITY FROM METRIC DIMENSIONAL REDUCTION AT SINGULARITIES

CRISTI STOICA

Abstract. A series of old and recent theoretical observations suggests that the quantization of gravity would be feasible, and some problems of Quantum Field Theory would go away if, somehow, the spacetime would undergo a dimensional reduction at high energy scales. But an identification of the cause of this dimensional reduction would still be desirable.

A possible explanation of the dimensional reduction is suggested by recent results in understanding the geometry of singularities in General Relativity. These new methods don't require modification of General Relativity, being just extensions of its mathematics to the limit cases. They turn out to work fine for some known types of cosmological singularities (black holes and FLRW Big-Bang), allowing a choice of the fundamental geometric invariants and physical quantities which remain regular. The resulting equations are equivalent to the standard ones outside the singularities.

One consequence of this mathematical approach to the singularities in General Relativity is a special, (geo)metric type of dimensional reduction: at singularities, the metric tensor becomes degenerate in certain spacetime directions, and some properties of the fields become independent of those directions. Effectively, it is like one or more dimensions of spacetime just vanish at singularities. Therefore, it seems that the geometry of singularities leads naturally to the spontaneous dimensional reduction needed by Quantum Gravity.

Contents

Introduction 2
1. Hints of dimensional reduction coming from other approaches 3
1.1. Dimensional reduction in Quantum Field Theory 3
1.2. Dimensional reduction in Quantum Gravity 7
2. Singularities in General Relativity 10
2.1. Benign singularities 10
2.2. Converting malign singularities into benign ones 12
3. Dimensional reduction at singularities 12
3.1. The dimension of the metric tensor 13
3.2. Metric dimension *vs.* topological dimension 13
3.3. Metric dimensional reduction and the Weyl tensor 15
3.4. Lorentz invariance and metric dimensional reduction 15
3.5. Particles lose two dimensions 16
3.6. Particles and spacetime anisotropy 16
3.7. The measure in the action integral 17
3.8. Does dimension vary with scale? 18
4. Conclusions 19
Acknowledgements 20
References 20

Introduction

Quantum Field Theory (QFT) and *General Relativity* (GR) are the most successful theories in fundamental theoretical physics. Their predictions were confirmed with very high precision, and they seem to offer accurate and complementary descriptions of the physical reality.

Yet, each one of them has some problems, especially when one tries to combine them. GR has the problems of infinities which appear at the *singularities.* QFT also has problems with infinities, which appear in the *perturbative expansion*, and are usually approached by renormalization techniques. Fortunately, the *renormalization group* formalized and solved many of the problems of QFT [1–6]. The *Standard Model* of particle physics is proven to be renormalizable [7–9], although the solution depends crucially on the existence of the Higgs boson.

But arguably the greatest difficulties appear when one tries to quantize gravity. General Relativity without matter fields is perturbatively non-renormalizable at two loops [10,11]. It requires an infinite number of higher derivative counterterms with their coupling constants. The main reason is the dimension of Newton's constant, which is $[\mathcal{G}_N] = 2 - D = -2$ in mass units.

In the quest of understanding the *small scale*, *ultraviolet* (UV) limit in QFT, and especially in the approaches to Quantum Gravity (QG), is accumulated evidence which seems to point in one particular direction. This evidence suggests, or even requires, that there is a *dimensional reduction* to two dimensions in the UV limit.

While many distinct approaches agree that somehow a dimensional reduction will solve the main problems of the interface between QFT and GR, what seems to be missing is the explicit cause leading to this spontaneous reduction.

As we will see in this paper, the dimensional reduction is ensured by the spacetime geometry at singularities in a very concrete way.

Usually, the apparent incompatibility between QFT and GR which manifests as non-renormalizability is considered to be the fault of the latter, hence usually the unification proposals start by modifying GR. Various approaches to QG are viewed as a hope which will cure not only the non-renormalizability, but also the problem of singularities, with the price of giving up one or more fundamental principles of GR.

Here we will take the opposite position: the solution to the problem of singularities comes from GR, and it also leads to the desired two-dimensionality in the UV limit, which is needed by quantum gravity.

The following can be considered a definition of Quantum Gravity (*cf.* 't Hooft [12]):

> Quantum Gravity is usually thought of as a theory, under construction, where the postulates of quantum mechanics are to be reconciled with those of general relativity, without allowing for any compromise in either of the two.

Here we will try to see how far we can go in reconciliating QFT and GR without making any compromises.

In this paper, we aim to show that the solution to the problems of singularities in GR, developed in [13–22] and reviewed briefly in §2, has implications to Quantum Gravity. Various approaches to QG suggest that if the spacetime becomes 2-dimensional at small scales, the quantization of gravity will become possible. Some of these hints will be reviewed in section §1. While dimensional reduction appears to be a desirable ingredient for QG, it would be useful to have an explanation of the reason which lead to the dimensional reduction, and a geometric interpretation of its meaning. In section §3, we will explain how the benign singularities cause the number of dimensions to be reduced, because of the way the metric becomes degenerate. Then we will try to connect the properties of the dimensional reduction caused by singularities, to those required by some of the approaches to QG.

1. **Hints of dimensional reduction coming from other approaches**

The method of regularization through dimensional reduction appeared from the observations that the loop integrations depend on the dimension in a continuous way, so that we can replace the dimension 4 by $4-\varepsilon$, avoiding the poles, and at the end make $\varepsilon \to 0$ [7, 8, 23]. The original method of dimensional regularization is rather formal, and apparently without implications to the actual physical dimensions. On the other hand, the fact that Quantum Gravity works fine in two dimensions justifies the consideration of the possibility that at small scales the number of dimensions is indeed reduced. In the following we review some "signs" that the spacetime is actually required to become two-dimensional in the small scale limit, while maintaining four dimensions at large scales.

1.1. **Dimensional reduction in Quantum Field Theory**

Suggestions that the two-dimensionality plays an important role appeared in various contexts of QFT. Since the first exactly solvable QFT model was discovered [24], the two-dimensional QFT proved to lead in a non-perturbative and direct manner to interesting results which can be applied then to make conjectures and find results for four-dimensions (see [25] and [26] and references therein).

1.1.1. **Two-dimensional QCD**

The two-dimensional theories are not necessarily just toy models. There are strong suggestions that the scattering amplitudes in QCD can be obtained from a two-dimensional field theory [27–29], and models in which "this two-dimensional nature of the interactions is manifest" appeared in the context of high energy Regge regime [30, 31].

1.1.2. **Fractal universe and measure dimensional reduction**

The *fractal universe program* developed by G. Calcagni originated from the idea of keeping from the Hořava-Lifschitz gravity the feature that it leads to a two-dimensional phase in the UV regime, but in the same time aims to remain Lorentz invariant. For this, it maintains an isotropic scaling, but to compensate it replaces the standard measure used in the action with a measure (initially a Lebesgue-Stieltjes measure) which reduces the Hausdorff dimension to two at the ultraviolet fixed point. The action becomes fractional, and the resulting theory is dissipative, hence non-unitary, although the energy turns out to be conserved.

The action is taken to be of the form

$$S=\int_{\mathcal{M}}\mathrm{d}\varrho(x)\,\mathcal{L}(\phi,\partial_{\mu}\phi) \tag{1}$$

with a measure

$$\mathrm{d}\varrho(x)=v\,\mathrm{d}x^{D} \tag{2}$$

Initially it was explored the theory with weight of the form $v:=\prod_{\mu=0}^{D-1}f_{(\mu)}(x)$. Taking all functions $f_{(\mu)}=f$ leads to an isotropic measure, but one can make also anisotropic choices.

As an example, the scalar field is described by the Lagrangian density

$$\mathcal{L}=-\frac{1}{2}\partial_{\mu}\phi\partial^{\mu}\phi-V(\phi) \tag{3}$$

where f is taken to have the dimension $[f]=1-\alpha$, so that $[v]=D(1-\alpha)$.

The scaling dimension of the scalar field ϕ is

$$[\phi]=\frac{D\alpha-2}{2} \tag{4}$$

and vanishes if and only if $\alpha=\frac{2}{D}$, or $\alpha=\frac{1}{2}$ for $D=4$.

If the potential is polynomial in ϕ

$$V=\sum_{n=0}^{N}\sigma_{n}\phi^{n}, \tag{5}$$

the resulting engineering dimension is

$$[\sigma_{N}]=D\alpha-\frac{N(D\alpha-2)}{2} \tag{6}$$

and to require the theory to be power-counting renormalizable, one imposes that N is restricted by the condition $[\sigma_N]\geq 0$. For $\alpha=\frac{D}{2}$, N is unconstrained, while for $D=4$ and $\alpha=1$, one can choose $N=4$. The idea is to construct a $D=4$ theory so that in the infrared limit $\alpha_{IR}=1$ and $N=4$, and in the ultraviolet limit, $\alpha_{UV}=\frac{1}{2}$.

Similar principles lead to a modified action for General Relativity, from which a modified version of Einstein's equation is derived.

The theory introduced in [32–34] was further refined and put on more rigorous mathematical basis in [35–41].

Mathematically, the fractal universe theory relied initially on the Lebesgue-Stieltjes measure (subsequently replaced by fractional measures [36]), fractional calculus, and fractional action principles [42–44].

In section §3.7 we will see that a measure of the form (2), having the desired properties, emerges naturally from our theory of singularities.

1.1.3. **Topological dimensional reduction**

The spacetime is considered, in general, to be a D-dimensional *topological manifold* – that is, its topology is locally like that of $\mathbb{R}^{D}$, and usually $D=4$. What would be the implications of a *topological dimensional reduction*, that is, if D would have, at some regions, a lower topological dimension?

In [45], D.V. Shirkov studied a $g\varphi^4$ QFT model described by a self-interacting Lagrangian $L = T - V$, where

$$V(m, g; \varphi) = \frac{m^2}{2}\varphi^2 + \frac{4\pi^{d/2}M^{4+d}}{9} g_d \varphi^4,\ g > 0 \tag{7}$$

This theory was initially motivated by the possible non-existence of a Higgs boson of 140±25, and aimed to explore the Ginzburg-Landau-Higgs alternative of a constant classical Higgs field at $\sim$ 250 GeV. But the absence of a Higgs quantum field makes the weak force again non-renormalizable. To obtain the regularization, D.V. Shirkov worked in a spacetime with variable topology, having a number of dimensions which varies from $D = 4$ in the IR limit, to $D = 1 + d < 4$ in the UV limit. The coupling constant was assumed to run from dimension $D = 4$ in the IR regime, to $D = 2$ in the UV regime.

One fundamental principle used was the *DR Agreement*, stipulating an equivalence between the reduction at spacetime scale $x_{dr} \sim \dfrac{1}{M_{dr}}$ and the reduction at the energy-momentum scale $p_{dr} \sim M_{dr}$.

The idea is illustrated by the approximation with a manifold obtained by joining two cylinders $S_{R,L}$ and $S_{r,l}$, of radii $R > r$ and lengths L, l, with a transition region S_{coll} of varying radius. Then, one can think at various problems taking place on the obtained manifold, for example solve some equations, and then take the limit $r \to 0$.

The replacement of the volume element in the momentum space

$$\mathrm{d}^4 k \to \mathrm{d}_M k = \frac{\mathrm{d}^4 k}{1 + \frac{k^2}{M^2}} \tag{8}$$

yields a one-loop Feynman integral which has as the IR asymptote the function

$$\ln \frac{q^2}{m_i^2}, \tag{9}$$

and as the UV asymptote the function

$$\ln \frac{4M^2}{m_i^2} + \frac{M^2}{q^2} \ln \frac{q^2}{M^2}. \tag{10}$$

This results in a reversal of the running coupling evolution in the two-dimensional region, and the value of the coupling constant gains a finite minimum value $\bar{g}_2(\infty) < \bar{g}_2(M_{dr}^2)$, where M_{dr} is the reduction scale. The coupling constant decreases in the IR limit as expected, but it has a maximum at $q = M_{dr}$, and in the UV limit decreases again, to the minimum $\bar{g}_2(\infty)$.

An interesting possibility which occurs is a novel type of *Grand Unified Theory* scenario, where the coupling constants of the Standard Model forces converge without needing an $SU(5)$ or other leptoquark symmetry [1]. As stated in the concluding section of [45]:

> The notable observation is that the change of geometry could yield the same final result as an explicit change of dynamics (by adding leptoquark fields etc.).

---

[1] For introductions to various GUT scenarios see [46, 47].

Of course, in this case the unification scale seems to require a bit higher energies than the GUT unification scale, taking place in the two-dimensional region, which is difficult to probe:

> Among further quests that are in order, let us put in the first place the issue of examining the chance of detecting some physical signal "through the looking-glass at scale M" that would provide us with direct evidence on the existence of dimension reduction of any kind.

The idea of topological dimensional reduction is further explored by P. Fiziev and D. V. Shirkov in [48–51], where it is applied to Klein-Gordon equation. In [48], the Klein-Gordon equation is studied on a spacetime which is the direct product between a 2-dimensional surface of revolution with variable radius and topology, and the 1-dimensional manifold $(\mathbb{R}^1, -\mathrm{d}t^2)$ representing the time dimension. The dimensional reduction takes place between $1+2$ and $1+1$ dimensions, by reducing the radius of revolution to 0.

By the method of separation of variables, the KG equation is reduced to three ordinary differential equations, from which two are simple wave equations. The third one is more difficult, but it can be put by a change of variables in the form of a one-dimensional Schrödinger-like equation, with the potential V defined by the geometry. The obtained solution forbids the propagation of signals related to the physical degrees of freedom related exclusively to the higher dimension, into the lower dimension region.

The resulting spectra depend on the mass from the KG equation and on the shape of the transition between the two regions of constant radii, and correspond to possible particles and their antiparticles. This suggests the idea that [48]

> The specific spectrum of scalar excitations resembles the spectrum of the real particles; it reflects the geometry of the transition region and represents its "fingerprints"

The possibility of deducing the geometry of the transition region by the "fingerprints" from the corresponding spectrum suggests an answer to the question about possible experimental "evidence on the existence of dimension reduction of any kind", asked in the first paper [45].

Some of these results were further generalized to higher dimension and multiple variable radii of compactification [49]. The manifold under consideration is a hypersurface in a semi-Euclidean $2d$-dimensional space $\mathbb{R}^{2d}_{2d-1}$ (see [52] for the notation), described by

$$\left\{\begin{array}{rlrl} x^0 = t, & \ldots, & x^{2k-1} = \rho_k(z)\cos(\phi_k), & \ldots, \\ x^{2d-1} = z, & \ldots, & x^{2k} = \rho_k(z)\sin(\phi_k), & \ldots \end{array}\right. \tag{11}$$

where $t \in \mathbb{R}$, $x \in \mathbb{R}$, $\phi_k \in [0, 2\pi]$, $\rho_k(z)$ are the shape functions (or the radii of compactification), and $k \in \{1, 2, \ldots, d-1\}$. This hypersurface is a Lorentzian manifold of dimension $1+d$. The solution turned out also in this general case to be reducible to one-dimensional Schrödinger-like equations, and to be regular even for $\rho_k = 0$.

The next step was to generalize the $(1+2)$-dimensional solution to the non-static case of *axial universes*, by allowing the shape function to depend also on time, $\rho = \rho(t, z) \geq 0$ [50]. A $(1+2)$-axial universe is a semi-Riemannian manifold which is a hypersurface of the Minkowski spacetime $\mathbb{R}^4_3$, defined by

$$\left\{\begin{array}{ll} x^0 = t, & x^1 = \rho(t, z)\cos(\phi), \\ x^3 = z, & x^2 = \rho(t, z)\sin(\phi) \end{array}\right. \tag{12}$$

where $t \in \mathbb{R}$, $x \in \mathbb{R}$, and $\phi \in [0, 2\pi]$.

The Einstein Equation translated into three equations connecting ρ and its partial derivatives with the energy-momentum tensor. Various energy-momentum tensors were considered in the Einstein Equation in $1+2$ dimensions:

1. in vacuum, where $\rho(t,z)$ was obtained to be linear in the variables t, z, respectively both of them,
2. with positive cosmological constant Λ, resulting in a static spherical universe,
3. in the presence of standard "dust", resulting in a homogeneous Monge-Ampère equation for ρ.

The solutions turned out to be integrable. The obtained dimension reduction points admit a classification, and there are hopes that they will provide new insights into the nature of the violations of the C, P, and T symmetries [48]:

> The parity violation due to the asymmetric character of the construction of our models could be related to violation of the CP symmetry.

Then, the Klein-Gordon equation was solved on the resulting spacetimes.

Section §3.2 establishes a strong connection between these results in topological dimensional reduction and our approach.

1.1.4. **Vanishing Dimensions**

One problem concerning the UV divergences in the Standard Model is the need to fine-tune the mass of the Higgs boson to an accuracy of 10^{-32}, to prevent the destabilization of the electroweak symmetry breaking scale [53]. Since fixing the cutoff at the electroweak scale makes the SM work well, it is believed that this is an indication of new physics beyond this scale. In [53] it is explored the alternative of keeping unmodified the structure of the SM, and reduce the dimension. One central point is that reducing the number of space dimensions to $d=2$ makes the Higgs terms linearly divergent, and $d=1$ makes them logarithmically divergent, making unnecessary the fine-tuning of the Higgs mass.

1.2. **Dimensional reduction in Quantum Gravity**

A review of the hints that, in various approaches to QG, a dimensional reduction occurs at small scales, is done by Carlip [54, 55]. Many of these hints involve the *spectral dimension*. The spectral dimension was calculated for causal dynamical triangulations (CDT) in [56], as evidence showing that the four-dimensional spacetime is recovered at larger scale. This resulted in a trend that various approaches to Quantum Gravity adhered to, consisting in calculating the spectral dimension in the UV limit to see if it is 2 [55, 57–59]. As it is known (see *e.g.* [41, 50, 60, 61]), while there is a correlation between the spectral dimension and the spacetime dimension, they are not equivalent. While the spectral dimension depends of the spacetime geometry too, it is very different. It represents the effective dimension of the diffusion process, being related to the dispersion relation of the corresponding differential operator. Spectral dimension is a widely used indicator in quantum geometry, and even if it is not equivalent with topological dimension, the latter bounds it.

As pointed out in [62], in dimension lower than four the Weyl curvature vanishes, and the vacuum Einstein equation has only locally flat solutions, or of constant curvature if the cosmological constant is not 0. This leads to the absence of local degrees of freedom, *i.e.* of gravitational waves for the Classical Gravity, and of gravitons for QG. Our own approach leads in a surprising way to this kind of dimensional reduction and vanishing of the Weyl curvature §3.3.

1.2.1. **The renormalization group: asymptotic safety**

General Relativity appears to be non-renormalizable, but the renormalization group analysis may give us useful hints. One possibility is that the infinite number of coupling constants become "unified" when approaching the UV fixed point. In [63] S. Weinberg proposed, as solution to the non-renormalizability of Quantum Gravity, the idea of *asymptotic safety.* The number of coupling constants needed by an asymptotically safe Quantum Gravity theory in four-dimensional spacetime is infinite, but their values are required to remain finite and to converge, under the renormalization group flow, to an ultraviolet fixed point. Also, the *ultraviolet critical surface* of the renormalization group flow is required to be finite-dimensional. In order to make the 3+1-dimensional gravity asymptotically safe, the Einstein-Hilbert Lagrangian density $\mathcal{L} = -\dfrac{1}{16\pi\mathcal{G}} R\sqrt{-\det g}$ has to be supplemented with higher-order curvature terms, for example by modifying it to $\mathcal{L} = f(R)\sqrt{-\det g}$. As stated in [63], "there is an asymptotically safe theory of pure gravity in $2+\varepsilon$ dimensions, with a one-dimensional critical surface", and "[a]symptotic safety is also preserved when we add matter fields", provided we add certain compensatory fields.

Some evidence accumulated in favor of asymptotic safety [64–69], especially near two dimensions [70, 71]. The spectral dimension near the fixed point appears to be $d_S = 2$ [57]. In [66] is showed that the existence of a non-Gaussian fixed point for the dimensionless coupling constant $g_N = \mathcal{G}_N \mu^{d-2}$ requires two-dimensionality.

1.2.2. **Causal dynamical triangulations**

In the *causal dynamical triangulations* approach [56, 72–75], spacetime is approximated by flat four-simplicial manifolds, similar to quantum Regge calculus [76]. The spacelike edges are taken to be of equal length, and the timelike edges of equal duration. The causality is enforced by requiring a fixed time-slicing at discrete times, and that the time-like edges agree in direction. The path integrals can be calculated non-perturbatively, resulting in four-dimensional spacetimes. The spectral dimension, which is the dimension as seen by a diffusion process, turns out to be four at large distances, but two at short distances [56].

1.2.3. **Hořava-Lifschitz gravity**

Inspired by the quantum critical phenomena in condensed matter systems, Hořava proposed in 2009 a model of Quantum Gravity [77]. The starting assumption is that the space and time behave differently at scaling – there is an anisotropic scaling invariance:

$$\begin{cases} \mathbf{x} & \mapsto b\mathbf{x}, \\ t & \mapsto b^z t. \end{cases} \tag{13}$$

To describe an UV fixed point, the critical exponent turns out to be $z = D - 1 = 3$, although it is argued that $z = 4$ would be even better. This anisotropy is not required to be a symmetry of the action itself, but of the solutions. The theory describes in the UV limit interacting non-relativistic gravitons, and is power-counting renormalizable in $1+3$ dimensions.

Lorentz invariance is absent in the UV limit, but it is conjectured that it emerges at large distances, where it is hoped that $z \to 1$. The resulting field equations are second-order in time, to avoid ghosts. In the same time they are of high order in space, canceling the divergences of the loop integrals. The spectral dimension turns out to be again two, for high energies, and four for low energies [58].

The anisotropy breaks the diffeomorphism invariance, and picks out a distinct time direction. This can be expressed in a space+time foliation $\mathcal{F}$, as in the ADM formalism [78]. The group of diffeomorphisms $\mathrm{Diff}(M)$ reduces to that of diffeomorphisms which preserve the leaves in the foliation, $\mathrm{Diff}_{\mathcal{F}}(M)$.

If the lapse function N depends on the time only, it is called *projectable*, otherwise it is called non-projectable. Theories with projectable N were the first to be studied, but now the hope moved toward the non-projectable ones.

The action splits naturally into a kinetic term $\mathcal{S}_K$ and a potential term $\mathcal{S}_V$:

$$\mathcal{S} = \mathcal{S}_K - \mathcal{S}_V. \tag{14}$$

The kinetic term $\mathcal{S}_K$ of the action is defined as the most general invariant term built from at most two time derivatives of the metric. In terms of the *extrinsic curvature*

$$K_{ij} := \frac{1}{2N}\left(\dot{g}_{ij} - \nabla_i N_j - \nabla_j N_i\right) \tag{15}$$

it has the form

$$\mathcal{S}_K = \frac{2}{\kappa^2}\int \left(K^{ij}K_{ij} - \lambda (K^s{}_s)^2\right)\sqrt{\det{}^3g}N\mathrm{d}t\mathrm{d}^3\mathrm{x} \tag{16}$$

where

$$\frac{2}{\kappa^2} = \frac{1}{16\pi\mathcal{G}_N}. \tag{17}$$

The dimension of the volume element is

$$[\mathrm{d}t\mathrm{d}^d\mathbf{x}] = -d - z \tag{18}$$

and each time derivative increases the dimension by

$$[\partial t] = z \tag{19}$$

therefore, the scaling dimension of κ is

$$[\kappa] = \frac{z-d}{2}, \tag{20}$$

so κ is dimensionless when $z = d$.

The potential term $\mathcal{S}_V$ includes terms up to six spatial derivatives of the metric.

The standard ADM action is

$$\mathcal{S}_{EH} = \frac{1}{16\pi\mathcal{G}_N}\int \left(K^{ij}K_{ij} - (K^s{}_s)^2 + {}^3R - 2\Lambda\right)\sqrt{\det{}^3g}N\mathrm{d}t\mathrm{d}^3\mathrm{x} \tag{21}$$

with the kinetic term $\lambda = 1$, which was hoped to emerge at large scale.

Some possible inconsistencies, internal and with the observations, in particular concerning the strong coupling and violations of unitarity, are discussed in [79–88]. Many of these objections arise from the difficulty to prove that GR is recovered in the IR limit.

A connection between our approach and Hořava-Lifschitz gravity is explored in section §3.6.

1.2.4. **Hints from other approaches**

As pointed out in [54, 55], there are hints from high temperature string theory [89] that the thermodynamic behavior becomes two-dimensional at high temperatures. Also, Modesto argued that in Loop Quantum Gravity the effective spectral dimension varies from four at large scales, to two at small scales [59, 90, 91]. Other results concern the spectral dimension in quantum spacetime based on noncommutative geometry [92, 93] and un-gravity [94].

2. Singularities in General Relativity

The other problem which seems to plague General Relativity is that of singularities. Under general conditions, the evolution equations in GR lead to singularities [95–100]. It seems that they are unavoidable. The options seem to be:

1. give up GR, or at least modify it, and
2. explore the singularities and try to find alternative but equivalent descriptions, which don't have problems with the infinities.

Since the first approach has been widely explored in the literature, we focused our research on the second one.

2.1. Benign singularities

Our initial intention was to make just a small step – to construct examples of singularities which can be worked out. We call these *benign singularities*. The main property they have is that the metric tensor g_{ab} is smooth, the singular features occurring because $\det g = 0$ at the singular points. This allows the construction of the Christoffel symbols of the first kind

$$\Gamma_{abc} := \frac{1}{2}\left(\partial_a g_{bc} + \partial_b g_{ca} - \partial_c g_{ab}\right). \tag{22}$$

But $\det g = 0$ forbids the construction of the Christoffel symbols of the second kind $\Gamma^{c}{}_{ab} := g^{cs}\Gamma_{abs}$, which involves the reciprocal metric $g^{ab} := (g^{-1})^{ab}$. Consequently, the curvature has to be defined in terms of the Christoffel symbols of the first kind, and not of the second, as it is normal:

$$R_{abcd} = \partial_a \Gamma_{bcd} - \partial_b \Gamma_{acd} + \Gamma_{ac\bullet}\Gamma_{bd\bullet} - \Gamma_{bc\bullet}\Gamma_{ad\bullet}. \tag{23}$$

The symbol $\bullet$ denote the contraction between covariant indices, and we adopted it because our contraction is more general than the usual one and the index notation would not be quite correct. Normally the contraction between covariant indices also requires the reciprocal metric g^{ab}, which becomes singular for $\det g = 0$. Luckily, the covariant contraction can be defined in a canonical and invariant way even in the case $\det g = 0$, provided that the tensor to be contracted satisfies certain conditions [13].

To see how, let's consider for starter two covectors $\omega, \tau \in T^*_p M$, where M is a semi-Riemannian manifold and $p \in M$. The contraction of the tensor $\omega \otimes \tau$ is given by $g^{st}\omega_s\tau_t$. This is defined by the metric g_{ab}, if $\det g \neq 0$. If $\det g = 0$, we can do this only if there are two vectors $u, v \in T_pM$ so that $\omega_a = g_{as}u^s$ and $\tau_a = g_{as}v^s$. In this case, we define the contraction by

$$\omega_\bullet \tau_\bullet := g_{st}u^s v^t. \tag{24}$$

We denote by $T_p{}^\bullet M$ the subspace of T_p^*M consisting of covectors (or 1-forms) of the form $\omega_a = (u^\flat)_a := g_{as}u^s$. We can extend this recipe to more general tensors from $\mathcal{T}^r_s(T_pM)$, provided that the components which we contract are from $T_p{}^\bullet M$. Note that we cannot use it to raise indices, since $u^\flat = (u+w)^\flat$ for any vector w so that $w^\flat = 0$. They form the kernel $\ker \flat$, which is zero if and only if $\det g = 0$.

From the above considerations we conclude that, to define the curvature as in (23), Γ_{abc} has to satisfy, at each point $p \in M$, the condition $\Gamma_{abs}w^s = 0$ for any vector $w^s \in T_pM$ which satisfies $g_{st}w^s v^t = 0$ for any vector $v \in T_pM$. Metrics satisfying this condition are named *radical-stationary*, and were studied by Kupeli for the case when the signature of the metric is constant [101, 102], and by the author for the general case [13, 14].

More details are given, in a manifestly invariant formulation, in [13], where we introduced also covariant derivatives for differential forms, and we defined the Riemann curvature tensor (23). The Riemann curvature tensor R_{abcd} can be defined in an invariant and canonical way, unlike $R^a{}_{bcd}$. Also, a simple condition ensured the smoothness of R_{abcd}, and such metrics and their singularities are named *semi-regular*. From the smoothness of R_{abcd} can be deduced in four dimensions the smoothness of $R_{ab}\det g$ and $R \det g$, and implicitly of

$$G_{ab}\det g := R_{ab}\det g - \frac{1}{2}g_{ab}R\det g. \tag{25}$$

This allows us to write a densitized version of Einstein's equation

$$G_{ab}\det g + \Lambda g_{ab}\det g = \kappa T_{ab}\det g, \tag{26}$$

where $\kappa := \dfrac{8\pi\mathcal{G}_N}{c^4}$. This equation is equivalent to Einstein's equation outside the singularities, but remains smooth at the singularities. The smoothness condition may be too strong sometimes, but what is important is that the metric is radical-stationary.

A condition stronger than semi-regularity is that of *quasi-regularity*, which allows a smooth Ricci decomposition of the Riemann tensor, and leads to another extension of Einstein's equation – the *expanded Einstein equation*, which is tensorial [20, 21]:

$$(G \circ g)_{abcd} + \Lambda (g \circ g)_{abcd} = \kappa (T \circ g)_{abcd} \tag{27}$$

where the operation

$$(h \circ k)_{abcd} := h_{ac}k_{bd} - h_{ad}k_{bc} + h_{bd}k_{ac} - h_{bc}k_{ad} \tag{28}$$

is the *Kulkarni-Nomizu product* of two symmetric bilinear forms h and k.

Big-Bang singularities of this type satisfy Penrose's Weyl curvature hypothesis [103, 104] automatically [22].

A simple but central result in singular semi-Riemannian geometry is the generalization, given in [14], of *warped products* [52]. The warped products allow the construction of a large class of semi-regular and quasi-regular singularities. As a corollary, a warped product of (non-degenerate) semi-Riemannian manifolds is semi-regular, and in some cases quasi-regular. The warped product can be used to show that the singularity at the Big-Bang of the Friedmann-Lemaître-Robertson-Walker model is semi-regular [19], and quasi-regular [20].

2.2. Converting malign singularities into benign ones

There are solutions to Einstein's equation for which one or more components of the metric tensor are singular – let's call them *malign singularities*. For example, the stationary black holes apparently are of this type. But they can be tamed, by a change of coordinates.

To understand how, let's recall that initially it was considered that the Schwarzschild solution has a singularity on the event horizon, in addition to that in the "center" of the black hole. This opinion lasted until Eddington proposed a coordinate system in which the metric became manifestly finite on the event horizon. The metric was in fact only apparently singular on the event horizon.

For the Schwarzschild [15], Reissner-Nordström [16], and Kerr-Newman [17] black holes there are coordinate changes which make the metric analytic at the genuine singularity $r = 0$. For example, the Reissner-Nordström solution

$$\mathrm{d}s^2 = -\left(1 - \frac{2m}{r} + \frac{q^2}{r^2}\right)\mathrm{d}t^2 + \left(1 - \frac{2m}{r} + \frac{q^2}{r^2}\right)^{-1}\mathrm{d}r^2 + r^2\mathrm{d}\sigma^2, \tag{29}$$

characterizing the stationary black holes with electric charge q and mass m, the component $g_{tt} \to \infty$ as $r \to 0$. A coordinate change of the form

$$\begin{cases} t &= \tau\rho^T \\ r &= \rho^S \end{cases} \tag{30}$$

transforms the metric to the form

$$\mathrm{d}s^2 = -\Delta\rho^{2T-2S-2}\left(\rho\mathrm{d}\tau + T\tau\mathrm{d}\rho\right)^2 + \frac{S^2}{\Delta}\rho^{4S-2}\mathrm{d}\rho^2 + \rho^{2S}\mathrm{d}\sigma^2, \tag{31}$$

where

$$\Delta := r^2 - 2mr + q^2 \text{ (hence } \Delta = \rho^{2S} - 2m\rho^S + q^2\text{)}. \tag{32}$$

For $T > S \geq 1$, the singularity turns out to be benign [16].

These coordinates suggest the possibility that the standard coordinate systems obtained for the black holes are in fact singular, and the correct coordinates are analytic, like those we have found.

In the new coordinates for the Reissner-Nordström and Kerr-Newman metrics, the electromagnetic potential and the electromagnetic fields also become analytic, and they are finite even for $r = 0$.

The new coordinates allow the spacetime to be foliated. In the Reissner-Nordström case, this is ensured by the condition $T \geq 3S$. By continuously modifying the parameters characterizing the black holes (m, q, and a), we can construct models in which they appear and disappear by Hawking evaporation. Because of the existence of foliations, we can construct globally hyperbolic spacetimes containing very general singularities [18]. Thus, the singularities don't necessarily destroy information.

3. Dimensional reduction at singularities

Adopting dimensional reduction only to justify Quantum Gravity would be an ad-hoc solution. Fortunately, the dimensional reduction emerges from the very structure of singularities.

This suggests the possibility that the solution to the problem of singularities helps understanding the problem of quantization of gravity.

3.1. The dimension of the metric tensor

If the metric tensor g_{ab} is degenerate at a point $p \in M$, then the distance in a part of the directions vanishes (fig. 1). The vanishing directions are given by the isotropic vectors from the vector subspace $\ker(\flat) := T_p^{\perp}(M) \leq T_pM$, consisting of the tangent vectors which are orthogonal to T_pM. These directions can be eliminated if we take the quotient space

$$T_{p\bullet}M := \frac{T_pM}{T_p^{\perp}(M)} \tag{33}$$

whose dimension is equal to the rank of the metric at p:

$$\dim T_{p\bullet}M = \operatorname{rank} g_p. \tag{34}$$

Then,

$$T_p{}^{\bullet}M := \flat(T_pM) \tag{35}$$

has the same dimension as $T_{p\bullet}M$, and in fact they are related by the isomorphism

$$\flat : T_{p\bullet}M \to T_p{}^{\bullet}M \tag{36}$$

induced by the morphism

$$\flat : T_pM \to T_p{}^{\bullet}M \leq T_p^*M. \tag{37}$$

We see that, because the metric tensor is degenerate, it can be reduced to a metric tensor on the space $T_{p\bullet}M$, and its inverse is a metric tensor on $T_p{}^{\bullet}M$. The dimension is actually given by the rank of the metric, which can be viewed as the dimension of the metric tensor.

3.2. Metric dimension *vs.* topological dimension

Let's consider a vector space V of dimension D. A symmetric bilinear form g on V defines a scalar product, or a metric. If g is degenerate, $\operatorname{rank} g < D$. The distance vanishes in the directions from $V^{\perp}$, and the geometric dimension is given by the rank of the metric. It is not necessarily equal to the vector space dimension D.

Let's consider now a semi-Riemannian manifold (M, g). Even if the rank of the metric is at some points lower than $D = \dim M$ (when the metric becomes degenerate), the topological dimension of the manifold remains D.

From mathematical viewpoint, in Differential Geometry there are three layers: the *topological structure*, the *differential structure* and the *geometric structure*. The topological structure on the set M is given by an atlas of local charts mapping an open set from M with one from $\mathbb{R}^D$, so that the transition maps are continuous. If the transition maps are differentiable, M becomes a differentiable manifold. If we add a metric tensor on M, we obtain a geometric structure. The topological dimension of M is the dimension $D = \dim \mathbb{R}^D$ of the vector space used in the charts of the atlas. The *metric dimension*, or the *geometric dimension* is given by the rank of the metric, and is allowed to be at most equal to the topological dimension.



FIGURE 1. (V, g) is an inner product vector space. The morphism $\flat : V \to V^*$ is defined by $u \mapsto u^\bullet := \flat(u) = u^\flat = g(u, \_)$. The radical $V_\circ := \ker \flat = V^\perp$ is the set of isotropic vectors in V. $V^\bullet := \operatorname{im} \flat \leq V^*$ is the image of $\flat$. The inner product g induces on $V^\bullet$ an inner product defined by $g_\bullet(u_1^\flat, u_1^\flat) := g(u_1, u_2)$, which is the inverse of g iff $\det g \neq 0$. The quotient $V_\bullet := V/V_\circ$ consists in the equivalence classes of the form $u+V_\circ$. On $V_\bullet$, g induces an inner product $g^\bullet(u_1+V_\circ, u_2+V_\circ) := g(u_1, u_2)$.

In the case when the metric is degenerate of constant signature (k, l, m) and is radical-stationary, a theorem of Kupeli [101] shows that the manifold (M, g) is locally isomorphic to a direct product manifold $P \times_0 N$ between a k-dimensional manifold N (without a metric, or with metric equal to 0) and a (non-degenerate) semi-Riemannian manifold P of signature (l, m). Hence, from the viewpoint of the metric g on M, at any point $p \in M$ the $k = D - \operatorname{rank} g$ dimensions associated with the degenerate directions can be ignored locally. We can thus identify the D-dimensional manifold M around the point p, with the rank g-dimensional manifold P. The manifold (M, g) looks locally like a lower-dimensional manifold (P, h). This situation is analogous to that of the gauge degrees of freedom.

If the metric is radical-stationary and has variable signature, the manifold (M, g) can be identified piecewisely with lower-dimensional manifolds (P, h). The information contained in the metric g of the manifold M can be obtained by pull-back from that of a metric on a manifold (P, h) with variable topological dimension.

This establishes the connection with the topological dimensional reduction proposed and studied by D.V. Shirkov and P. Fiziev (see section §1.1.3). Our approach leads to a conclusion which is very close to the following observation from [50]:

> dimensional reduction of the physical space in general relativity (GR) can be regarded as an unrealized and as yet untapped consequence of Einstein's equations (EEqs) themselves which takes place around singular points of their solutions.

It is not clear at this point to what extent the geometric reduction of radical-stationary semi-Riemannian manifolds is equivalent to the semi-Riemannian manifolds with variable topological dimension. But what we can say is that in GR there are other fields to be considered in addition to the metric tensor. The information contained in those fields will be, in general, lost by the topological reduction induced by the metric dimensional reduction. On the other hand, in order to admit smooth covariant contractions and smooth covariant derivatives (as defined in [13] for differential forms and tensors with covariant indices), the fields are required to ignore to some extent the degenerate directions. But even under these conditions, they are more general than the fields defined on manifolds of lower topological dimension, and we cannot recover the former from the latter ones by pull-back.

Keeping the points topologically distinct, even though the distance induced by the metric vanishes between them, provides more generality than allowing the topological dimensional reduction.

One important reason to avoid making the topological identification due to the dimensional reduction is that variable topological dimension is not compatible with the foliation of the spacetime in spacelike hypersurfaces. This kind of foliation is important for global hyperbolicity and for avoiding the information loss [18].

3.3. Metric dimensional reduction and the Weyl tensor

At quasi-regular singularities the Ricci decomposition is smooth. According to a theorem we proved in [22], the Weyl curvature vanishes at quasi-regular singularities. The reason is that $\dim T_p{}^{\bullet}M < D$, hence the dimension of the Riemann curvature tensor R_{abcd} is reduced. For $D = 4$, $\dim T_p{}^{\bullet}M \leq 3$, consequently any tensor having the algebraic properties of the Weyl tensor vanishes. Because the Ricci decomposition is smooth, in particular the Weyl curvature tensor C_{abcd} is smooth, and this means that around the singularity the Weyl tensor remains small.

An example of quasi-regular singularity is the Schwarzschild black hole [21], which can be used as a classical model for neutral spinless particles. At this time it is not clear whether the singularities of the stationary charged and rotating black holes are quasi-regular, but they are analytic, the geometric dimensional reduction occurs, and the electromagnetic potential and its field are analytic and finite at $r = 0$ [16, 17].

The vanishing of the Weyl curvature tensor implies that the local degrees of freedom, *i.e.* the gravitational waves for GR and the gravitons for QG, are absent [62]. Because of continuity, as approaching a quasi-regular singularity, the contribution of the gravitons vanishes, and this diminishes the renormalizability problems of QG.

3.4. Lorentz invariance and metric dimensional reduction

When the metric of a spacetime becomes degenerate, it is no longer Lorentzian. The group of transformations which is associated to the metric can't be a Lorentz group, since the metric is degenerate. The role of the Lorentz group is taken by a larger group – the *Barbilian group* [105]. But if the metric is radical-stationary, the Barbilian group can be reduced to a subgroup, which is a Lorentz group of lower dimension.

We can say, because of this, that the condition to be radical-stationary is the mathematical expression of the Lorentz invariance, in the case when the metric is allowed to be degenerate.

The spacetimes with this kind of metric satisfy the Lorentz invariance so long as the metric is non-degenerate, and when it becomes degenerate, the Lorentz invariance is maintained, but for lower dimensions. One should mention here that this is the best we can do, if we want to include the singularities in the spacetime. Having the full 4-dimensional Lorentz invariance at singularities is not possible, because of the very definition of singularities.

By comparison, other approaches to QG had to give up Lorentz invariance even outside the singularities. It is the case of loop quantum gravity and Hořava-Lifschitz gravity for example, where it is hoped that the four-dimensional Lorentz invariance emerges at large scales.

3.5. **Particles lose two dimensions**

As described in [16, 17] and in §2.2, a charged black hole of type Reissner-Nordström or Kerr-Newman can be described by an analytic metric, and the electromagnetic potential and its field are also analytic and remain finite at $r = 0$. From the viewpoint of GR, charged particles are such black holes. Also this applies to other types of gauge fields, such as the Yang-Mills fields[2].

As we can see from (31), in our coordinates, at $\rho = 0$ (which is equivalent to $r = 0$), the metric loses two dimensions, those corresponding to coordinates ρ and τ. Apparently the metric on the sphere vanishes too, but this only reflects that the warped product describing spherically symmetric solutions to Einstein's equation involves all the concentric spheres, down to $r = 0$.

Metric dimensional reduction occurs similarly for the Schwarzschild (describing neutral particles) and the more general Kerr-Newman (describing charged or neutral, spinning or not particles) cases.

The fact that two dimensions are lost, and that the gauge potential and fields remain finite at the singularity, are expected to have important impact to field quantization.

One of the major problems of electrodynamics is the fact that the particle's potential and field become infinite as approaching $r = 0$. This problem turns out to be removed, by employing our non-singular coordinate system. This sets as one priority in the future developments of our program to see exactly what happens in the perturbative expansions and in the renormalization group analysis.

3.6. **Particles and spacetime anisotropy**

The metric (31) admits a foliation in spacelike hypersurfaces only for $T \geq 3S$ in (30) [16]. This condition allows the coordinates to be compatible with the distinction between space and time. But it leads to an anisotropy between space and time, which is manifest when passing to the old Reissner-Nordström coordinates (t, r). A rescaling in the coordinates (τ, ρ) is isotropic, of course in the coordinates (τ, ρ), but the coordinates (t, r) are rescaled anisotropically, due to (30).

The diffeomorphism invariance should be considered valid in coordinates (τ, ρ), but not in the singular coordinates (t, r).

This anisotropic scaling invariance is similar to that which P. Hořava managed to obtain by modifying the Lagrangian of GR in [77]. His analysis shows that the anisotropy would lead to the correct dimension for the Newton constant (see §1.2.3 and [77]). Apparently, the anisotropy we obtained in [16] is not equivalent to that from Hořava-Lifschitz gravity, because ours follows from standard GR considerations, while that of Hořava from modifications of the Einstein-Hilbert Lagrangian (and implicitly of Einstein's equation).

---

[2] Of course, the relativistic effects should be complemented by the quantum properties to get a complete description of a physical particle, and we don't have yet such a model.

In our proposal, the metric is still the fundamental field. The very degeneracy of the metric imposes conditions on the foliation at the singularity. There is no need for other structure to define the foliation, as it is in Hořava-Lifschitz gravity. In our approach there is no need to impose in the IR limit the recovery of standard GR, since we start from standard GR outside the singularities, and extend it in the singularities.

3.7. **The measure in the action integral**

When the metric becomes degenerate, its determinant vanishes. Consequently, the volume form

$$\mathrm{d}_{vol} := \sqrt{-\det g}\mathrm{d}x^0 \wedge \ldots \wedge \mathrm{d}x^{D-1} = \sqrt{-\det g}\mathrm{d}x^D \tag{38}$$

tends to 0 as approaching the degenerate singularities (in non-singular coordinates, in which the metric is smooth).

The action principle is given by

$$S = \int_{\mathcal{M}} \mathrm{d}_{vol}(x)\, \mathcal{L} \tag{39}$$

If the metric is diagonal in the coordinates (x^μ), then

$$\mathrm{d}_{vol}(x) = \prod_{\mu=0}^{D-1} \sqrt{|g_{\mu\mu}(x)|}\mathrm{d}x^D. \tag{40}$$

This is in fact similar to the measure studied by G. Calcagni [32, 33]. This is obvious if we take in the Lebesgue measure from §1.1.2, (1), the following weights:

$$f_{(\mu)}(x) = \sqrt{|g_{\mu\mu}(x)|}. \tag{41}$$

This identification makes much of the analysis developed in [32,33] for QFT be a consequence of the fact that the metric may be degenerate.

In terms of $v(x) = \prod_{\mu=0}^{D-1} f_{(\mu)}(x)$, we have

$$v = \sqrt{-\det g}, \tag{42}$$

so that the measure becomes

$$\mathrm{d}\varrho(x) = \sqrt{-\det g}\mathrm{d}x^D, \tag{43}$$

which is just the standard measure from General Relativity, except that in our framework it is allowed to vanish.

The justification for changing the measure in G. Calcagni's proposal comes in his theory from the hypothesis that the spacetime is fractal, the Hausdorff dimension changing with the scale. The fractal nature manifests in the usage of fractional calculus. He uses a general Lebesgue-Stieltjes measure, independent of GR, and applies it to flat spacetime QFT, as well as to QG [32–34].

The action (1) refers to the Special Relativistic QFT, but the identification (41) we proposed replaces the weigts f_μ with the square root of the metric components, which we allowed to tend to zero. This suggests that GR comes to rescue QFT at high energies, by reducing the dimension to two. For GR, Calcagni applies the same recipe he used in Special Relativistic QFT: proposes that the weight v multiplies $\sqrt{-\det g}$ [32]. This would of course lead to a modified Einstein equation, and modified GR. Our approach sticks to the standard GR, the only difference being in allowing the metric to become degenerate.

In our solution, the measure is simply due to the degeneracy of the metric. Our claim is that GR itself provides the measure and tames the infinities of QFT.

3.8. **Does dimension vary with scale?**

So far we provided arguments that the singularities characterized by the degeneracy of the metric explain the geometric dimensional reduction. This dimensional reduction has many common features with the dimensional reduction expected in other proposals in the literature. In addition, it is not invented with the problem of quantization in mind, but it is a consequence of our approach to the singularity problem, which in turn fits naturally in classical GR.

Yet, the hardest part remains to be done. The geometric dimensional reduction we propose becomes manifest as the distance to a singularity becomes smaller. But the dimensional reduction needed in QFT and QG seems to have nothing to do with the distance to a singularity. It is required just to depend on the scale.

The precise ways in which the geometric dimensional reduction we proposed impacts QFT and GR need to be analyzed in more depth. At this point, we will give only a "qualitative" (*i.e.* "handwaving") justification.

The higher order Feynman diagrams involve a larger number of particles. This means that in the same region of space there will be more particles, which we will consider to be benign singularities. Because of this, the metric will have, in average, smaller determinant. Recall that for benign singularities the determinant of the metric tends to 0 as the distance to the singularity decreases, and having a higher number of singularities in the same region reduces the average of the metric's determinant (fig. 2). Thus, in the high energy limit, the measure dimensional reduction will become more and more present in the integrals.

Let's discuss a bit the diffeomorphism invariance (general covariance) of the above argument. It is clear that both $\det g$ and other relevant quantities such as the components of the Weyl tensor C_{abcd} depend on the coordinates. The singularities describing the particles provide some constraints that $\det g = 0$, and $C_{abcd} = 0$ for quasi-regular singularities, along the universe lines describing the particles. These constraints are invariant under local diffeomorphisms. It is true that changing the coordinates one can make these quantities increase away from the singularities, but the constraints are invariant. Also, $\det g$ is present in the action, and the action integral is invariant.

In conclusion, *the main conjecture* is that, although the dimensional reduction happens at singularities which represent the particles, when many particles are present, the measure dimensionality – present through $\det g$ – is reduced in average. We claim that this acts like a regulator. Let's name this *the hypothesis of average dimensional reduction*.

If this hypothesis is true, then the fact that the average metric determinant changes with the scale is not an absolute law, but merely an "accident". If the real reason of change is



FIGURE 2. Schematic picture illustrating how we expect that the metric's average determinant decreases as the number of singularities (*i.e.* particles) in the region increases. The red dots represent the singularities, and the blue line represents $|\det g|$.

the way the singularities are distributed in a given region, then this may have observable consequences at larger scales too. But at larger scales, the predictions of GR we expected so far seem to be confirmed with high degree of accuracy, yet nothing like this has been found. On the other hand, let's not forget that most tests of GR concern phenomena involving a small number of bodies which are separated by large regions of vacuum. We need to derive and test predictions for the case when matter is distributed more homogeneously, as the galaxies appear at large scale. And we know that at large scale, the galaxies don't seem to fit what we presently expect to be the behavior predicted by GR, as we understand it so far. It would be interesting to check if some of the discrepancies usually attributed to *dark matter* can in fact be explained as large scale manifestations of the average dimensional reduction. This may be supported by the idea that the galaxies rotate as if they were $(1+2)$-dimensional [106].

4. **Conclusions**

We reviewed some of the hints indicating that if a dimensional reduction would take place at small scales, then some major problems concerning the quantization of gravity, but also of other fields, would go away. Some hints refer to the dimension involved in calculations, others to the geometric and topological dimensions, and others to the spectral dimension.

We advocated here the position that the approach of singularities introduced and developed in [13–22], leading to a (geo)metric dimensional reduction, also lays a foundation for the quantization of gravity. This position is supported by the strong connections between the metric dimensional reduction and the other kinds of dimensional reductions, reviewed in this paper.

This is just a small step; many questions remain open, and much work remains to be done.

Acknowledgements

The author thanks P. Fiziev and D. V. Shirkov for helpful discussions and advice received during his stay at the Bogoliubov Laboratory of Theoretical Physics, JINR, Dubna.

References

[1] ECG Stueckelberg and A. Petermann. Normalization of the constants in the theory of quanta. *Helvetica Physica Acta (Switzerland)*, 26, 1953.

[2] M. Gell-Mann and FE Low. Quantum electrodynamics at small distances. *Physical Review*, 95(5):1300, 1954.

[3] N.N. Bogoliubov and D.V. Shirkov. *Nuovo Cim.*, 3:845–863, 1956.

[4] N.N. Bogoliubov and D.V. Shirkov. *Introduction to the theory of quantized fields*. John Wiley & Sons, 1980.

[5] D.V. Shirkov. The Bogoliubov Renormalization Group. *arXiv:hep-th/9602024*, 1996. arXiv:hep-th/9602024.

[6] D.V. Shirkov. The Bogoluibov Renormalization Group in Theoretical and Mathematical Physics . *arXiv:hep-th/9903073*, 1999.

[7] G. 't Hooft and M. Veltman. Regularization and renormalization of gauge fields. *Nuclear Physics B*, 44(1):189–213, 1972.

[8] G. 't Hooft. Dimensional regularization and the renormalization group. *Nuclear Physics B*, 61:455–468, 1973.

[9] G. Hooft. The glorious days of physics-renormalization of gauge theories. *arXiv:hep-th/9812203*, 1998.

[10] G. 't Hooft and M. Veltman. One loop divergencies in the theory of gravitation. In *Annales de l'Institut Henri Poincaré: Section A, Physique théorique*, volume 20, pages 69–94. Institut Henri Poincaré, 1974.

[11] M.H. Goroff and A. Sagnotti. The ultraviolet behavior of Einstein gravity. *Nuclear Physics B*, 266(3-4):709–736, 1986.

[12] Gerard 't Hooft. The fundamental nature of space and time. In Daniele Oriti, editor, *Approaches to Quantum Gravity: Toward a New Understanding of Space, Time and Matter*, pages 13–36. Cambridge University Press, 2009. Online version.

[13] C. Stoica. On Singular Semi-Riemannian Manifolds. *arXiv:math.DG/1105.0201*, May 2011.

[14] C. Stoica. Warped Products of Singular Semi-Riemannian Manifolds. *arXiv:math.DG/1105.3404*, May 2011.

[15] C. Stoica. Schwarzschild Singularity is Semi-Regularizable. *arXiv:gr-qc/1111.4837*, November 2011.

[16] C. Stoica. Analytic Reissner-Nordstrom Singularity. *Phys. Scr.*, 85:055004, November 2012. arXiv:gr-qc/1111.4332.

[17] C. Stoica. Kerr-Newman Solutions with Analytic Singularity and no Closed Timelike Curves. *arXiv:gr-qc/1111.7082*, November 2011.

[18] C. Stoica. Globally Hyperbolic Spacetimes with Singularities. *arXiv:math.DG/1108.5099*, August 2011.

[19] C. Stoica. Big Bang singularity in the Friedmann-Lemaitre-Robertson-Walker spacetime. *arXiv:gr-qc/1112.4508*, December 2011.

[20] C. Stoica. Beyond the Friedmann-Lemaitre-Robertson-Walker Big Bang singularity. *arXiv:gr-qc/1203.1819*, March 2012.

[21] C. Stoica. Einstein equation at singularities. *arXiv:gr-qc/1203.2140*, March 2012.

[22] C. Stoica. On the Weyl Curvature Hypothesis. *arXiv:gr-qc/1203.3382*, March 2012.

[23] CG Bollini and JJ Giambiagi. Dimensional renorinalization: The number of dimensions as a regularizing parameter. *Il Nuovo Cimento B (1971-1996)*, 12(1):20–26, 1972.

[24] W.E. Thirring. A soluble relativistic field theory. *Annals of Physics*, 3(1):91–112, 1958.

[25] E. Abdalla, M.C.B. Abdalla, and K.D. Rothe. *Non-perturbative methods in 2 dimensional quantum field theory*. World Scientific Pub Co Inc, 1991.

[26] Y. Frishman and J. Sonnenschein. *Non-perturbative Field Theory: From Two-dimensional Conformal Field Theory to QCD in Four Dimensions*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2010.

[27] LN Lipatov. Massless particle bremsstrahlung theorems for high-energy hadron interactions. *Nuclear Physics B*, 307(4):705–720, 1988.

[28] LN Lipatov. Review in Perturbative QCD, ed. AH Mueller, 1989.

[29] LN Lipatov. High-energy scattering in QCD and in quantum gravity and two-dimensional field theories. *Nuclear Physics B*, 365(3):614–632, 1991.

[30] H. Verlinde and E. Verlinde. QCD at high energies and two-dimensional field theory. *arXiv:hep-th/9302104*, 1993.

[31] I.Y. Aref'eva. Regge regime in QCD and asymmetric lattice gauge theory. *Physics Letters B*, 325(1):171–182, 1994.

[32] G. Calcagni. Quantum field theory, gravity and cosmology in a fractal universe. *Journal of High Energy Physics*, 2010(3):1–38, 2010. arXiv:hep-th/1001.0571.

[33] G. Calcagni. Fractal universe and quantum gravity. *Physical review letters*, 104(25):251301, 2010. arXiv:hep-th/0912.3142.

[34] G. Calcagni. Gravity on a multifractal. *Physics Letters B*, 2011. arXiv:hep-th/1012.1244.

[35] G. Calcagni. Discrete to continuum transition in multifractal spacetimes. *Physical Review D*, 84(6):061501, 2011. arXiv:hep-th/1106.0295.

[36] G. Calcagni. Geometry of fractional spaces. *arXiv:hep-th/1106.5787*, 2011.

[37] G. Calcagni. Geometry and field theory in multi-fractional spacetime. *Journal of High Energy Physics*, 2012(1):1–77, 2012. arXiv:hep-th/1107.5041.

[38] M. Arzano, G. Calcagni, D. Oriti, and M. Scalisi. Fractional and noncommutative spacetimes. *Physical Review D*, 84(12):125002, 2011. arXiv:hep-th/1107.5308.

[39] G. Calcagni and G. Nardelli. Momentum transforms and laplacians in fractional spaces. *arXiv:hep-th/1202.5383*, 2012.

[40] G. Calcagni. Diffusion in quantum gravity. *arXiv:hep-th/1204.2550*, 2012.

[41] G. Calcagni. Diffusion in multi-fractional spacetimes. *arXiv:hep-th/1205.5046*, 2012.

[42] R.A. El-Nabulsi. A fractional action-like variational approach of some classical, quantum and geometrical dynamics. *International Journal of Applied Mathematics*, 17(3):299, 2005.

[43] R.A. El-Nabulsi and D.F.M. Torres. Fractional actionlike variational problems. *Journal of Mathematical Physics*, 49(5):053521–053521, 2008.

[44] C. Udrişte and D. Opriş. Euler-Lagrange-Hamilton dynamics with fractional action. *WSEAS Transactions on Mathematics*, 7(1):19–30, 2008.

[45] D.V. Shirkov. Coupling running through the looking-glass of dimensional reduction. *Phys. Part. Nucl. Lett.*, 7(6):379–383, 2010. arXiv:hep-th/1004.1510.

[46] H. Georgi. *Lie algebras in particle physics: From isospin to unified theories*. Benjamin/Cummings Pub. Co., Advanced Book Program (Reading, Mass.), 1982.

[47] Baez, J. And Huerta, J. The Algebra of Grand Unified Theories. *American Mathematical Society*, 47(3):483–552, 2010.

[48] P.P. Fiziev and D.V. Shirkov. Solutions of the Klein-Gordon equation on manifolds with variable geometry including dimensional reduction. *Theoretical and Mathematical Physics*, 167(2):680–691, 2011. arXiv:hep-th/1009.5309.

[49] P.P. Fiziev. Riemannian (1+d)-Dim Space-Time Manifolds with Nonstandard Topology which Admit Dimensional Reduction to Any Lower Dimension and Transformation of the Klein-Gordon Equation to the 1-Dim Schrödinger Like Equation. *arXiv:math-ph/1012.3520*, 2010.

[50] P.P. Fiziev and D.V. Shirkov. The (2+1)-dim Axial Universes – Solutions to the Einstein Equations, Dimensional Reduction Points, and Klein–Fock–Gordon Waves. *J. Phys. A*, 45(055205):1–15, 2012. arXiv:gr-qc/arXiv:1104.0903.

[51] D.V. Shirkov. Dream-land with Classic Higgs field, Dimensional Reduction and all that. In *Proceedings of the Steklov Institute of Mathematics*, volume 272, pages 216–222, 2011.

[52] B. O'Neill. Semi-Riemannian Geometry with Applications to Relativity. *Pure Appl. Math.*, (103):468, 1983.

[53] L. Anchordoqui, D.C. Dai, M. Fairbairn, G. Landsberg, and D. Stojkovic. Vanishing dimensions and planar events at the LHC. *arXiv:hep-ph/1003.5914*, 2010.

[54] S. Carlip, J. Kowalski-Glikman, R. Durka, and M. Szczachor. Spontaneous dimensional reduction in short-distance quantum gravity? In *Aip Conference Proceedings*, volume 31, page 72, 2009.

[55] S. Carlip. The Small Scale Structure of Spacetime. *arXiv:gr-qc/1009.1136*, 2010.

[56] J. Ambjørn, J. Jurkiewicz, and R. Loll. Spectral dimension of the universe. *Physical review letters*, 95(17):171301, 2005. arXiv:hep-th/0505113.

[57] O. Lauscher and M. Reuter. Fractal spacetime structure in asymptotically safe gravity. *Journal of High Energy Physics*, 10:050, 2005.

[58] P. Hořava. Spectral dimension of the universe in quantum gravity at a lifshitz point. *Physical review letters*, 102(16):161301, 2009.

[59] L. Modesto. Fractal structure of loop quantum gravity. *Class.Quant.Grav*, 26:242002, 2009. arXiv:gr-qc/0812.2214.

[60] T.P. Sotiriou, M. Visser, and S. Weinfurtner. Spectral dimension as a probe of the ultraviolet continuum regime of causal dynamical triangulations. *Physical Review Letters*, 107(13):131303, 2011.

[61] T.P. Sotiriou, M. Visser, and S. Weinfurtner. From dispersion relations to spectral dimension?nd back again. *Physical Review D*, 84(10):104018, 2011.

[62] S. Carlip. Lectures in (2+ 1)-dimensional gravity. *J.Korean Phys.Soc*, 28:S447–S467, 1995. arXiv:gr-qc/9503024.

[63] S. Weinberg. Ultraviolet divergences in quantum theories of gravitation. In *General relativity: an Einstein centenary survey*, volume 1, pages 790–831, 1979.

[64] M. Reuter and F. Saueressig. Renormalization group flow of quantum gravity in the Einstein-Hilbert truncation. *Physical Review D*, 65:065016, 2002. arXiv:hep-th/0110054.

[65] D.F. Litim. Fixed points of quantum gravity. *Physical review letters*, 92(20):201301, 2004. arXiv:hep-th/0312324.

[66] M. Niedermaier. The asymptotic safety scenario in quantum gravity: An introduction. *Classical and Quantum Gravity*, 24:R171, 2007. arXiv:gr-qc/0610018.

[67] H.W. Hamber and R.M. Williams. Nonlocal effective gravitational field equations and the running of newton? constant g. *Physical Review D*, 72(4):044026, 2005. arXiv:hep-th/0507017.

[68] M. Reuter and F. Saueressig. Functional renormalization group equations, asymptotic safety, and quantum Einstein gravity. *arXiv:hep-th/0708.1317*, 2007.

[69] A. Codello, R. Percacci, and C. Rahmede. Investigating the ultraviolet properties of gravity with a Wilsonian renormalization group equation. *Annals of Physics*, 324(2):414–469, 2009. arXiv:hep-th/0805.2909.

[70] H. Kawai, Y. Kitazawa, and M. Ninorniya. Renormalizability of quantum gravity near two dimensions. *Nuclear Physics B*, 467(1):313–331, 1996. arXiv:hep-th/9511217.

[71] D. Litim. On fixed points of quantum gravity. In *A Century of relativity physics: ERE 2005, XXVIII Spanish Relativity Meeting, Oviedo, Asturias, Spain, 6-10 September 2005*, volume 841, pages 322–329. Amer Inst of Physics, 2006.

[72] J. Ambjørn, J. Jurkiewicz, and R. Loll. Nonperturbative Lorentzian path integral for gravity. *Physical Review Letters*, 85(5):924–927, 2000. arXiv:hep-th/0002050.

[73] J. Ambjørn, J. Jurkiewicz, and R. Loll. Emergence of a 4D world from causal quantum gravity. *Physical Review Letters*, 93(13):131301, 2004. arXiv:hep-th/0404156.

[74] J. Ambjørn, J. Jurkiewicz, and R. Loll. Reconstructing the universe. *Physical Review D*, 72(6):064014, 2005. arXiv:hep-th/0505154.

[75] J. Ambjørn, J. Jurkiewicz, and R. Loll. Quantum gravity, or the art of building spacetime. In Daniele Oriti, editor, *Approaches to Quantum Gravity: Toward a New Understanding of Space, Time and Matter*, pages 341–359. Cambridge University Press, 2009. arXiv:hep-th/0604212.

[76] T. Regge. Nuovo cimento 19 558 collins pa and williams rm 1974. *Phys. Rev. D*, 10:3537, 1961.

[77] P. Hořava. Quantum Gravity at a Lifshitz Point. *Physical Review D*, 79(8):084008, 2009. arXiv:hep-th/0901.3775.

[78] R. Arnowitt, S. Deser, and C.W. Misner. The Dynamics of General Relativity, in: Gravitation: An Introduction to Current Research. pages 227–?64, 1962.

[79] C. Charmousis, G. Niz, A. Padilla, and P.M. Saffin. Strong coupling in hořava gravity. *Journal of High Energy Physics*, 2009:070, 2009.

[80] S. Weinfurtner, T.P. Sotiriou, and M. Visser. Projectable Hořava–lifshitz gravity in a nutshell. In *Journal of Physics: Conference Series*, volume 222, page 012054. IOP Publishing, 2010.

[81] T.P. Sotiriou. Hořava-lifshitz gravity: a status report. In *Journal of Physics: Conference Series*, volume 283, page 012034. IOP Publishing, 2011.

[82] M. Visser. Status of Hořava gravity: A personal perspective. In *Journal of Physics: Conference Series*, volume 314, page 012002. IOP Publishing, 2011.

[83] D. Blas, O. Pujolas, and S. Sibiryakov. On the extra mode and inconsistency of hořava gravity. *Journal of High Energy Physics*, 2009:029, 2009.

[84] I. Kimpton and A. Padilla. Lessons from the decoupling limit of hořava gravity. *Journal of High Energy Physics*, 2010(7):1–26, 2010.

[85] M. Henneaux, A. Kleinschmidt, and G.L. Gómez. A dynamical inconsistency of hořava gravity. *Physical Review D*, 81(6):064002, 2010.

[86] A. Papazoglou and T.P. Sotiriou. Strong coupling in extended horava-lifshitz gravity. *Physics Letters B*, 685(2-3):197–200, 2010.

[87] D. Blas, O. Pujolas, and S. Sibiryakov. Comment on ?trong coupling in extended hořava–lifshitz gravity?phys. lett. b 685 (2010) 197]. *Physics Letters B*, 688(4):350–355, 2010.

[88] A. Wang and Q. Wu. Stability of spin-0 graviton and strong coupling in horava-lifshitz theory of gravity. *Physical Review D*, 83(4):044025, 2011.

[89] J.J. Atick and E. Witten. The hagedorn transition and the number of degrees of freedom of string theory. *Nuclear Physics B*, 310(2):291–334, 1988.

[90] L. Modesto. Fractal quantum space-time. *arXiv:gr-qc/0905.1665*, May 2009.

[91] F. Caravelli and L. Modesto. Fractal dimension in 3d spin-foams. *Arxiv preprint arXiv:0905.2170*, May 2009.

[92] D. Benedetti. Fractal properties of quantum spacetime. *Physical review letters*, 102(11):111303, 2009.

[93] L. Modesto and P. Nicolini. Spectral dimension of a quantum universe. *Physical Review D*, 81(10):104040, 2010.

[94] P. Nicolini and E. Spallucci. Un-spectral dimension and quantum spacetime phases. *Physics Letters B*, 695(1):290–293, 2011.

[95] R. Penrose. Gravitational Collapse and Space-Time Singularities. *Phys. Rev. Lett.*, (14):57–59, 1965.

[96] S. Hawking. The occurrence of singularities in cosmology. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 294(1439):511–521, 1966.

[97] S. Hawking. The occurrence of singularities in cosmology. ii. *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, pages 490–493, 1966.

[98] S. Hawking. The occurrence of singularities in cosmology. iii. causality and singularities. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 300(1461):187–201, 1967.

[99] S. Hawking and R. Penrose. The Singularities of Gravitational Collapse and Cosmology. *Proc. Roy. Soc. London Ser. A*, (314):529–548, 1970.

[100] S. Hawking and G. Ellis. *The Large Scale Structure of Space Time*. Cambridge University Press, 1995.

[101] D. Kupeli. Degenerate Manifolds. *Geom. Dedicata*, 23(3):259–290, 1987.

[102] D. Kupeli. *Singular Semi-Riemannian Geometry*. Kluwer Academic Publishers Group, 1996.

[103] R. Penrose. Singularities and time-asymmetry. In *General relativity: an Einstein centenary survey*, volume 1, pages 581–638, 1979.

[104] R. Penrose. *Cycles of time: an extraordinary new view of the universe*. Alfred a Knopf Inc, 2011.

[105] Dan Barbilian. Galileische Gruppen und quadratische Algebren. *Bull. Math. Soc. Roumaine Sci.*, page XLI, 1939.

[106] W.F. Kao and R.T. Yu. The Effect of the (2+1)-Dimensional Gravity in the Spiral Galaxy. *arXiv:astro-ph/0008268*, August 2000.